Author Index Vol. 434

Adam, M., see Macek, J. 423
Allain, P., see Premel-Cabic, A. 187
Anjaneyulu, B., see Bhatia, S.C. 288
Aronsson, E., see Odham, G. 31
Assmann, B.
— and Haas, H.J.
Determination of the age dependency of the creatinine-related pseudouridine excretion in children's urine by high-performance liquid chromatography 202
Babušíková, O., see Boháček, J. 439
Balíková, M.
— and Kohlíček, J.
Rapid determination of ethylene glycol at toxic levels in serum and urine 469
Bamonte, F., see Dionisotti, S. 169
Banavalikar, M.M., see Bhatia, S.C. 288
Bargnoux, H., see Bougerolle, A.M. 232
Baveja, S.K., see Padmalatha Devi, K. 265
Belenkii, B.G.
Review. Use of laser detectors in capillary liquid chromatography 337
Berger, J.A., see Bougerolle, A.M. 232
Bergqvist, Y.
— and Churchill, F.C.
Review. Detection and determination of antimalarial drugs and their metabolites in body fluids 1
Bharucha, E.D., see Bhatia, S.C. 296, 303
Bhatia, K.L., see Kamar, S. 228
Bhatia, S.C.
—, Revankar, S.N., Anjaneyulu, B., Desai, N.D., Banavalikar, M.M., Sahajwalla, C.F. and Subrahmanyam, D.
High performance liquid chromatography of antifilarials. I. Simultaneous determination of 1-4-(4-nitrophenylaminophenylthiocarbamido-4-methyl)piperazine and three of its metabolites in biological fluids 288
—, Revankar, S.N., Bharucha, E.D., Desai, N.D. and Sahajwalla, C.G.
High-performance liquid chromatography of antifilarials. III. Determination of 2-*tert.*-butyl-6-isothiocyanato-5-(piperidin-1-yl)-benzthiazole in biological fluids 303
—, Revankar, S.N., Doshi, K.J., Desai, N.D., Bharucha, E.D. and Sahajwalla, C.G.
High-performance liquid chromatography of antifilarials. II. Determination of S-(2-carboxyethyl)-N-(2-*tert.*-butyl-5-methoxybenzthiazol-6-yl) dithiocarbamate and its active isothiocyanate metabolite in biological fluids 296
Bidault, R., see Le Coguic, A. 320
Bisse, E.
— and Wieland, H.
High-performance liquid chromatographic separation of human haemoglobins. Simultaneous quantitation of foetal and glycated haemoglobins 95
Bökens, H.
—, Bourscheidt, C. and Müller, R.F.
Determination of bumetanide in plasma by high-performance liquid chromatography 327
Boháček, J.
—, Hošek, B., Babušíková, O. and Hrivňáková, A.
Determination of adenosine deaminase and purine nucleoside phosphorylase activities using high-performance liquid chromatographic and radiochromatographic methods 439
Bougerolle, A.M.
—, Chabard, J.L., Jbilou, M., Bargnoux, H., Petit, J., Berger, J.A. and Dordain, G.
Simultaneous determination of trimipramine and its demethylated metabolites in plasma by gas chromatography–mass spectrometry 232
Bourscheidt, C., see Bökens, H. 327
Caccamo, F.
—, Carfagnini, G., Di Corcia, A. and Samperi, R.
Measurement of urinary estriol glucuronides during the menstrual cycle by high-performance liquid chromatography 61
Carfagnini, G., see Caccamo, F. 61
Černý, M., see Jegorov, A. 417
Chabard, J.L., see Bougerolle, A.M. 232
Churchill, F.C., see Bergqvist, Y. 1
Clark, B.J., see Fell, A.F. 377
Clarke, G.S., see Sheridan, M.E. 253
Clifford, A.J., see Smidt, C.R. 21
Coguic, A. Le, see Le Coguic, A. 320
Cook, C.E., see Nakahara, Y. 247
Corcia, A. Di, see Caccamo, F. 61
Cserháti, T.
— and Szögyi, M.
Amino acid–amino acid interactions studied by charge-transfer chromatography 455

Curtis, M.A., see Pullen, R.H. 271
Dauphin, A., see Le Coguic, A. 320
Dayer, P., see Padmalatha Devi, K. 265
Desai, N.D., see Bhatia, S.C. 288, 296, 303
Devi, K. Padmalatha, see Padmalatha Devi, K. 265
Di Corcia, A., see Caccamo, F. 61
Dionisotti, S.
—, Bamonte, F., Gamba, M. and Ongini, E.
High-performance liquid chromatographic determination of netilmicin in guinea-pig and human serum by fluorodinitrobenzene derivatization with spectrophotometric detection 169
Doormaal, J.J. van, see Hoving, E.B. 395
Dordain, G., see Bougerolle, A.M. 232
Doshi, K.J., see Bhatia, S.C. 296
Dubiel, W., see Toguzov, R.T. 447
Dubois, J.P., see Sioufi, A. 239
Ehrsson, H., see Hassan, M. 330
Farinotti, R., see Le Coguic, A. 320
Fell, A.F.
—, Noctor, T.A.G., Mama, J.E. and Clark, B.J.
Computer-aided optimisation of drug enantiomer separation in chiral high-performance liquid chromatography 377
Frič, P., see Kocna, P. 429
Gamba, M., see Dionisotti, S. 169
Gerber, G., see Toguzov, R.T. 447
Gibassier, D., see Sado, P.A. 157
Haas, H.J., see Assmann, B. 202
Hack, M.H.
— and Helmy, F.M.
Application of methylamine O-deacylation and Vitride reduction to the analysis of neutral lipids: a thin-layer chromatographic evaluation 215
Hais, I.M., see Nobilis, M. 363
Hase, S.
—, Ikenaka, K., Mikoshiba, K. and Ikenaka, T.
Analysis of tissue glycoprotein sugar chains by two-dimensional high-performance liquid chromatographic mapping 51
Hassan, M.
— and Ehrsson, H.
Determination of busulfan in human plasma by gas chromatography with electron-capture detection 330
Hayashi, S., see Matsukura, H. 196
Hekkens, W.Th.J.M., see Kocna, P. 429
Helmy, F.M., see Hack, M.H. 215
Hill, H.M., see Selinger, K. 259
Hirata, N.
—, Kasai, M., Yanagihara, Y. and Noguchi, K.
Separation of peptides by multimode chromatography on a column packed with vinyl alcohol copolymer gel 71
Holthuis, J.J.M., see Vendrig, D.E.M.M. 145
Hoogmartens, J., see Kibwage, I.O. 177
Hošek, B., see Boháček, J. 439
Hoving, E.B.
—, Jansen, G., Volmer, M., Van Doormaal, J.J. and Muskiet, F.A.J.
Profiling of plasma cholesterol ester and triglyceride fatty acids as their methyl esters by capillary gas chromatography, preceded by a rapid aminopropyl-silica column chromatographic separation of lipid classes 395
—, Prins, J., Rutgers, H.M. and Muskiet, F.A.J.
Behaviour of plasmalogens during high-performance liquid chromatography on a silica column with a mobile phase containing phosphoric acid 411
Hrivňáková, A., see Boháček, J. 439
Ichinose, H., see Naoi, M. 209
Ikenaka, K., see Hase, S. 51
Ikenaka, T., see Hase, S. 51
Jansen, G., see Hoving, E.B. 395
Jbilou, M., see Bougerolle, A.M. 232
Jegorov, A.
—, Tříska, J., Trnka, T. and Černý, M.
Separation of α-amino acid enantiomers by reversed-phase high-performance liquid chromatography after derivatization with *o*-phthaldialdehyde and a sodium salt of 1-thio-β-D-glucose 417
Jones, A.D., see Smidt, C.R. 21
Kaiser, D.G.
—, Vangiessen, G.J., Shah, J.A. and Weber, D.J.
Gas chromatographic determination of guanadrel in plasma and urine 135
Kaiser, G., see Sioufi, A. 239
Kamar, S.
— and Bhatia, K.L.
Characterization of buffalo lactotransferrin by polyacrylamide gel electrophoresis in 6 *M* urea 228
Karnes, H.T., see Sarkar, M.A. 324
Kasafírek, E., see Kocna, P. 429
Kasai, M., see Hirata, N. 71
Kennedy, C.M., see Pullen, R.H. 271
Kibwage, I.O.
—, Roets, E., Verbruggen, A., Hoogmartens, J. and Vanderhaeghe, H.
Thin-layer chromatographic study of the metabolites of erythromycins in the Wistar rat 177

Kocna, P.
—, Frič, P., Kočová-Holáková, M., Slabý, J., Kasafírek, E. and Hekkens, W.Th. J.M.
Isolation and analysis of peptidic fragments of α-gliadin using reversed-phase high-performance liquid chromatography 429
Kočová-Holáková, M., see Kocna, P. 429
Koel, M., see Nebinger, P. 43
Kohlíček, J., see Balíková, M. 469
Kurachi, M., see Kurata, K. 278
Kurata, K.
—, Kurachi, M. and Tanii, Y.
High-performance liquid chromatographic determination of mianserin in plasma and brain and its application to pharmacokinetic studies in the rat 278
LaCreta, F.P.
—, Olszewski, J.J. and Tew, K.D.
Purification of glutathione S-transferases from rat liver and Walker 256 mammary carcinoma cells by high-performance liquid chromatography and a glutathione affinity column 83
Lamiable, D.
—, Vistelle, R., Nguyen-Khac, M. and Millart, H.
High-performance liquid chromatographic determination of dexamethasone in cerebrospinal fluid and plasma in the rabbit 315
Le Coguic, A.
—, Bidault, R., Farinotti, R. and Dauphin, A.
Determination of ofloxacin in plasma and urine by liquid chromatography 320
Leemann, T., see Padmalatha Devi, K. 265
Leverge, R., see Sado, P.A. 157
Lichý, A., see Macek, J. 423
McDowall, M., see Odham, G. 31
Macek, J.
—, Lichý, A., Tesařová, E. and Adam, M.
Separation of type IX collagen from other cartilage collagens by hydrophobic interaction chromatography 423
Mama, J.E., see Fell, A.F. 377
Maniwa, M., see Umeno, Y. 123
Marunaka, T., see Umeno, Y. 123
Matheou, D., see Selinger, K. 259
Matsukura, H.
—, Suzuki, Y., Okada, T., Naiki, S., Hayashi, S. and Yamaji, Y.
High-performance liquid chromatographic determination of acid and alkaline phosphatases in urine 196
Matsushima, E., see Umeno, Y. 123
Meiborg, G., see Volmer, M. 385
Mezzetti, T.
—, Proietti, G. and Sbordoni, A.
Simple method for the determination of hormonal iodine and exogenous iodine from X-ray contrast media drugs 308
Michelsen, P., see Odham, G. 31
Mikoshiba, K., see Hase, S. 51
Millart, H., see Lamiable, D. 315
Muda, M.
—, Pelizzoni, F., Sello, G. and Mussini, E.
Determination of ergothioneine in red blood cells by high-performance liquid chromatography 191
Müller, R.F., see Bökens, H. 327
Muskiet, F.A.J., see Hoving, E.B. 395, 411
—, see Volmer, M. 385
Mussini, E., see Muda, M. 191
Nagatsu, T., see Naoi, M. 209
Naiki, S., see Matsukura, H. 196
Nakahara, Y.
— and Cook, C.E.
Confirmation of cannabis use. III. Simultaneous quantitation of six metabolites of Δ^9-tetrahydrocannabinol in plasma by high-performance liquid chromatography with electrochemical detection 247
Naoi, M.
—, Ichinose, H., Takahashi, T. and Nagatsu, T.
Sensitive assay for determination of pyridoxal-5-phosphate in enzymes using high-performance liquid chromatography after derivatization with cyanide 209
Nebinger, P.
— and Koel, M.
Automated analysis of urinary 3-methoxy-4-hydroxymandelic acid using ion-pair chromatography and fluorimetric detection 43
Nguyen-Khac, M., see Lamiable, D. 315
Nobilis, M.
— and Hais, I.M.
Direct absorptiometric quantitation in high-performance liquid chromatography in the absence of concomitant reference standards 363
Noctor, T.A.G., see Fell, A.F. 377
Noguchi, K., see Hirata, N. 71
Odham, G.
—, Valeur, A., Michelsen, P., Aronsson, E. and McDowall, M.
Highly sensitive determination and characterization of intact cellular ester-linked phospholipids using liquid chromatography–plasma spray mass spectrometry 31

Okada, T., see Matsukura, H. 196
Olivares, J.
— and Verdys, M.
Isocratic high-performance liquid chromatographic method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 111
Olszewski, J.J., see LaCreta, F.P. 83
Ongini, E., see Dionisotti, S. 169
Padmalatha Devi, K.
—, Ranga Rao, K.V., Baveja, S.K., Leemann, T. and Dayer, P.
Determination of alprenolol and metoprolol in plasma by column liquid chromatography 265
Pelizzoni, F., see Muda, M. 191
Peteghem, C. Van, see Rosier, J. 222
Petit, J., see Bougerolle, A.M. 232
Pimenov, A.M., see Toguzov, R.T. 447
Pommier, F., see Sioufi, A. 239
Premel-Cabic, A.
— and Allain, P.
Simultaneous liquid chromatographic determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 187
Prins, J., see Hoving, E.B. 411
Proietti, G., see Mezzetti, T. 308
Prokudin, V.Yu., see Toguzov, R.T. 447
Pullen, R.H.
—, Kennedy, C.M. and Curtis, M.A.
Direct plasma injection using internal surface reversed-phase high-performance liquid chromatography: feasibility study using propofol as a model compound 271
Ranga Rao, K.V., see Padmalatha Devi, K. 265
Rao, K.V. Ranga, see Padmalatha Devi, K. 265
Revankar, S.N., see Bhatia, S.C. 288, 296, 303
Richter, K.
Determination of clozapine in human serum by capillary gas chromatography 465
Roets, E., see Kibwage, I.O. 177
Rosier, J.
— and Van Peteghem, C.
Detection of oxidatively modified 2′-deoxyguanosine-3′-monophosphate, using ^{32}P-postlabelling and anion-exchange thin-layer chromatography 222
Rutgers, H.M., see Hoving, E.B. 411
Sado, P.A.
—, Gibassier, D. and Leverge, R.
Dosage simultané de la vinburnine et de la 6-hydroxyvinburnine par chromatographie liquide à haute performance 157
Sahajwalla, C.G., see Bhatia, S.C. 288, 296, 303
Samperi, R., see Caccamo, F. 61
Sarkar, M.A.
— and Karnes, H.T.
Comparison of solid-phase extraction and liquid–liquid extraction for high-performance liquid chromatographic analysis of antipyrine in plasma 324
Sbordoni, A., see Mezzetti, T. 308
Selinger, K.
—, Matheou, D. and Hill, H.M.
High-performance liquid chromatographic method for the determination of miconazole in vaginal fluid 259
Sello, G., see Muda, M. 191
Shah, J.A., see Kaiser, D.G. 135
Sheridan, M.E.
— and Clarke, G.S.
Improved high-performance liquid chromatographic determination of doxycycline in serum and urine using solid-phase extraction columns 253
Shimizu, T., see Shioya, Y. 283
Shioya, Y.
— and Shimizu, T.
High-performance liquid chromatographic procedure for the determination of a new anti-gastric ulcer agent, 2-(4-chlorobenzoylamino)-3-[2(1H)-quinolinon-4-yl]propionic acid, in human plasma and urine 283
Sioufi, A.
—, Pommier, F., Kaiser, G. and Dubois, J.P.
Determination of benazepril, a new angiotensin-converting enzyme inhibitor, and its active metabolite, benazeprilat, in plasma and urine by capillary gas chromatography–mass-selective detection 239
Slabý, J., see Kocna, P. 429
Smidt, C.R.
—, Jones, A.D. and Clifford, A.J.
Gas chromatography of retinol and α-tocopherol without derivatization 21
Subrahmanyam, D., see Bhatia, S.C. 288
Suzuki, Y., see Matsukura, H. 196
Szögyi, M., see Cserháti, T. 455
Takahashi, T., see Naoi, M. 209
Tanii, Y., see Kurata, K. 278
Teeuwsen, J., see Vendrig, D.E.M.M. 145
Tesařová, E., see Macek, J. 423
Tew, K.D., see LaCreta, F.P. 83

Tikhonov, Y.V., see Toguzov, R.T. 447
Toguzov, R.T.
—, Tikhonov, Y.V., Pimenov, A.M., Prokuding, V.Yu., Dubiel, W., Ziegler, M. and Gerber, G.
Optimization of the ion-pair high-performance liquid chromatographic separation of purine derivatives in erythrocytes, thymocytes and liver mitochondria 447
Tříska, J., see Jegorov, A. 417
Trnka, T., see Jegorov, A. 417
Umeno, Y.
—, Matsushima, E., Maniwa, M. and Marunaka, T.
Gas chromatographic–negative-ion chemical ionization mass spectrometric determination of a new dihydropyridine calcium antagonist (MPC-1304) and its metabolites in human plasma and urine 123
Valeur, A., see Odham, G. 31
Vanderhaege, H., see Kibwage, I.O. 177
Van Doormaal, J.J., see Hoving, E.B. 395
Vangiessen, G.J., see Kaiser, D.G. 135
Van Peteghem, C., see Rosier, J. 222
Vendrig, D.E.M.M.
—, Teeuwsen, J. and Holthuis, J.J.M.
Determination of vinca alkaloids in plasma and urine using ion-exchange chromatography on silica gel and fluorescence detection 145
Verbruggen, A., see Kibwage, I.O. 177
Verdys, M., see Olivares, J. 111
Vistelle, R., see Lamiable, D. 315
Volmer, M., see Hoving, E.B. 395
—, Meiborg, G. and Muskiet, F.A.J.
Simultaneous capillary gas chromatographic profiling of medium- and long-chain fatty acid methyl esters with split injection. Correction for injection-related discrimination by the 'bracketing' method 385
Weber, D.J., see Kaiser, D.G. 135
Wieland, H., see Bisse, E. 95
Yamaji, Y., see Matsukura, H. 196
Yanagihara, Y., see Hirata, N. 71
Ziegler, M., see Toguzov, R.T. 447

Subject Index Vol. 434

Absorptiometric quantitation
Direct absorptiometric quantitation in HPLC in the absence of concomitant reference standards 363

Acid phosphatase
HPLC determination of acid and alkaline phosphatases in urine 196

Adenosine deaminase
Determination of adenosine deaminase and purine nucleoside phosphorylase activities using HPLC and radiochromatographic methods 439

Alkaline phosphatase
HPLC determination of acid and alkaline phosphatases in urine 196

Alprenolol
Determination of alprenolol and metoprolol in plasma by column LC 265

Amino acid–amino interactions
Amino acid–amino acid interactions studies by charge-transfer chromatography 455

α-Amino acid enantiomers
Separation of α-amino acid enantiomers by reversed-phase HPLC after derivatization with *o*-phthaldialdehyde and a sodium salt of 1-thio-β-D-glucose 417

Amino acids
Amino acid–amino acid interactions studies by charge-transfer chromatography 455

Amodiaquine
Review. Detection and determination of antimalarial drugs and their metabolites in body fluids 1

Anhydroerythromycin A
TLC study of the metabolites of erythromycins in the Wistar rat 177

Antifilarials
HPLC of antifilarials. I. Simultaneous determination of 1-4-(4-nitrophenylaminophenylthiocarbamido-4-methyl)piperazine and three of its metabolites in biological fluids 288

Antifilarials
HPLC of antifilarials. II. Determination of S-(2-carboxyethyl)-N-(2-*tert.*-butyl-5-methoxybenzthiazol-6-yl) dithiocarbamate and its active isothiocyanate metabolite in biological fluids 296

Antifilarials
HPLC of antifilarials. III. Determination of 2-*tert.*-butyl-6-isothiocyanato-5-(piperidin-1-yl)benzthiazole in biological fluids 303

Antimalarial drugs
Review. Detection and determination of antimalarial drugs and their metabolites in body fluids 1

Antipyrine
Comparison of solid-phase extraction and liquid–liquid extraction for HPLC analysis of antipyrine in plasma 324

Benazepril
Determination of benazepril, a new angiotensin-converting enzyme inhibitor, and its active metabolite, benazeprilat, in plasma and urine by capillary GC–mass-selective detection 239

Benazeprilat
Determination of benazepril, a new angiotensin-converting enzyme inhibitor, and its active metabolite, benazeprilat, in plasma and urine by capillary GC–mass-selective detection 239

Benflurene
Direct absorptiometric quantitation in HPLC in the absence of concomitant reference standards 363

Benzodiazepines
Computer-aided optimisation of drug enantiomer separation in chiral HPLC 377

Bumetanide
Determination of bumetanide in plasma by HPLC 327

Busulfan
Determination of busulfan in human plasma by GC with electron-capture detection 330

2-*tert.*-Butyl-6-isothiocyanato-5-(piperidin-1-yl)benzthiazole
HPLC of antifilarials. III. Determination of 2-*tert.*-butyl-6-isothiocyanato-5-(piperidin-1-yl)benzthiazole in biological fluids 303

S-(2-Carboxyethyl)-N-(2-*tert.*-butyl-5-methoxybenzthiazol-6-yl) dithiocarbamate
HPLC of antifilarials. II. Determination of S-(2-carboxyethyl)-N-(2-*tert.*-butyl-5-methoxybenzthiazol-6-yl) dithiocarbamate and its active isothiocyanate metabolite in biological fluids 296

Catecholamines
Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 187
Chiral stationary phases, LC
Computer-aided optimisation of drug enantiomer separation in chiral HPLC 377
2-(4-Chlorobenzoylamino)-3-[2(1H)-quinolinon-4-yl]propionic acid
HPLC procedure for the determination of a new anti-gastric ulcer agent, 2-(4-chlorobenzoylamino)-3-[2(1H)-quinolinon)-4-yl]propionic acid, in human plasma and urine 283
Chloroquine
Review. Detection and determination of antimalarial drugs and their metabolites in body fluids 1
Cholesterol ester
Profiling of plasma cholesterol ester and triglyceride fatty acids as their methyl esters by capillary GC, preceded by a rapid aminopropyl-silica column chromatographic separation of lipid classes 395
Clozapine
Determination of clozapine in human serum by capillary GC 465
Collagen type IX
Separation of type IX collagen from other cartilage collagens by hydrophobic interaction chromatography 423
Cytidine
Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 111
Cytosine
Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 111
N-Demethylerythromycin A
TLC study of the metabolites of erythromycins in the Wistar rat 177
Deoxycytidine
Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 111
2′-Deoxyguanosine-3′-monophosphate
Detection of oxidatively modified 2′-deoxyguanosine-3′-monophosphate, using ^{32}P-postlabelling and anion-exchange TLC 222
Dexamethasone
HPLC determination of dexamethasone in cerebrospinal fluid and plasma in the rabbit 315
Dihydropyridine calcium antagonist
GC–negative-ion chemical ionization MS determination of new dihydropyridine calcium antagonist (MPC 1304) and its metabolites in human plasma and urine 123
3,4-Dihydroxyphenylacetic acid
Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 187
3,4-Dihydroxyphenylalanine
Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 187
3,4-Dihydroxyphenylethyleneglycol
Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethyleneglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 187
Dihydroxy-Δ^9-tetrahydrocannabinols
Confirmation of cannabis use. III. Simultaneous quantitation of six metabolites of Δ^9-tetrahydrocannabinol in plasma by HPLC with electrochemical detection 247
Doxycycline
Improved HPLC determination of doxycycline in serum and urine using solid-phase extraction columns 253
Drugs
Computer-aided optimisation of drug enantiomer separation in chiral HPLC 377
Enantiomer separation
Computer-aided optimisation of drug enantiomer separation in chiral HPLC 377
Enzymes
Purification of glutathione S-transferases from rat liver and Walker 256 mammary cells by HPLC and a glutathione affinity column 83
Enzymes
HPLC determination of acid and alkaline phosphatases in urine 196
Enzymes
Sensitive assay for determination of pyridoxal-5-phosphate in enzymes using HPLC after derivatization with cyanide 209

Enzymes
Determination of adenosine deaminase and purine nucleoside phosphorylase activities using HPLC and radiochromatographic methods 439
Ergothioneine
Determination of ergothioneine in red blood cells by HPLC 191
Erythromycins
TLC study of the metabolites of erythromycins in the Wistar rat 177
Estriol glucuronides
Measurement of urinary estriol glucuronides during the menstrual cycle by HPLC 61
Ethylene glycol
Rapid determination of ethylene glycol at toxic levels in serum and urine 469
Fatty acid methyl esters
Simultaneous capillary GC profiling of medium- and long-chain fatty acid methyl esters with split injection correction for injection-related discrimination by the 'bracketing' method 385
Fatty acids
Profiling of plasma cholesterol ester and triglyceride fatty acids as their methyl esters by capillary GC, preceded by a rapid aminopropyl-silica column chromatographic separation of lipid classes 395
Fluorescence detectors
Review. Use of laser detectors in capillary LC 337
Foetal haemoglobins
HPLC separation of human haemoglobins. Simultaneous quantitation of foetal and glycated haemoglobins 95
α-Gliadin
Isolation and analysis of peptidic fragments of α-gliadin using reversed-phase HPLC 429
Glutathione S-transferases
Purification of glutathione S-transferases from rat liver and Walker 256 mammary carcinoma cells by HPLC and a glutathione affinity column 83
Glycated haemoglobins
HPLC separation of human haemoglobins. Simultaneous quantitation of foetal and glycated haemoglobins 95
Glycoprotein sugar chains
Analysis of tissue glycoprotein sugar chains by two-dimensional HPLC 51
Guanadrel
GC determination of guanadrel in plasma and urine 135
Haemoglobins
HPLC separation of human haemoglobins. Simultaneous quantitation of foetal and glycated haemoglobins 95
11-Hydroxy-Δ^9-tetrahydrocannabinol
Confirmation of cannabis use. III. Simultaneous quantitation of six metabolites of Δ^9-tetrahydrocannabinol in plasma by HPLC with electrochemical detection 247
6-Hydroxyvinburnine
Simultaneous determination of vinburnine and 6-hydroxyvinburnine in human plasma by HPLC 157
Iodine
Simple method for the determination of hormonal iodine and exogenous iodine from X-ray contrast media drugs 308
Lactotransferrin
Characterization of buffalo lactotransferrin by polyacrylamide gel electrophoresis in 6 *M* urea 228
Laser detectors
Review. Use of laser detectors in capillary LC 337
Lipids
Application of methylamine O-deacylation and Vitride reduction to the analysis of neutral lipids: a TLC evaluation 215
Lipids
Profiling of plasma cholesterol ester and triglyceride fatty acids as their methyl esters by capillary GC, preceded by a rapid aminopropyl-silica column chromatographic separation of lipid classes 395
Long-chain fatty acid methyl esters
Simultaneous capillary GC profiling of medium- and long-chain fatty acid methyl esters with split injection. Correction for injection-related discrimination by the 'bracketing' method 385
Lorazepam
Computer-aided optimisation of drug enantiomer separation in chiral HPLC 377
Medium-chain fatty acid methyl esters
Simultaneous capillary GC profiling of medium- and long-chain fatty acid methyl esters with split injection correction for injection-related discrimination by the 'bracketing' method 385
Mefloquine
Review. Detection and determination of antimalarial drugs and their metabolites in body fluids 1

3-Methoxy-4-hydroxymandelic acid
Automated analysis of urinary 3-methoxy-4-hydroxymandelic acid using ion-pair chromatography and fluorimetric detection 43
Methylamine O-deacylation
Application of methylamine O-deacylation and Vitride reduction to the analysis of neutral lipids: a TLC evaluation 215
(±)-Methyl 2-oxopropyl 1,4-dihydro-2,6-dimethyl-4-(2-nitrophenyl)-3,5-pyridinedicarboxylate
GC–negative-ion chemical ionization MS determination of a new dihydropyridine calcium antagonist (MPC 1304) and its metabolites in human plasma and urine 123
Metoprolol
Determination of alprenolol and metoprolol in plasma by column LC 265
Mianserin
HPLC determination of mianserin in plasma and brain and its application to pharmacokinetic studies in the rat 278
Miconazole
HPLC method for the determination of miconazole in vaginal fluid 259
Netilmicin
HPLC determination of netilmicin in guinea-pig and human serum by fluorodinitrobenzene derivatization with spectrophotometric detection 169
1-4-(4-Nitrophenylaminophenylthiocarbamido-4-methyl) piperazine
HPLC of antifilarials. I. Simultaneous determination of 1-4-(4-nitrophenylaminophenylthiocarbamido-4-methyl)piperazine and three of its metabolites in biological fluids 288
Norepinephrine
Simultaneous LC determination of norepinephrine, 3,4-dihydroxyphenylethyleglycol, 3,4-dihydroxyphenylalanine and 3,4-dihydroxyphenylacetic acid in human plasma 187
11-Nor-Δ^9-tetrahydrocannabinol-9-COOH
Confirmation of cannabis use. III. Simultaneous quantitation of six metabolites of Δ^9-tetrahydrocannabinol in plasma by HPLC with electrochemical detection 247
Nucleobases
Optimization of the ion-pair HPLC separation of purine derivatives in erythrocytes, thymocytes and liver mitochondria 447
Nucleosides
Optimization of the ion-pair HPLC separation of purine derivatives in erythrocytes, thymocytes and liver mitochondria 447
Nucleotides
Optimization of the ion-pair HPLC separation of purine derivatives in erythrocytes, thymocytes and liver mitochondria 447
Ofloxacin
Determination of ofloxacin in plasma and urine by LC 320
Oligosaccharides
Analysis of tissue glycoprotein sugar chains by two-dimensional HPLC 51
Oxamniquine
Computer-aided optimisation of drug enantiomer separation in chiral HPLC 377
Oxazepam
Computer-aided optimisation of drug enantiomer separation in chiral HPLC 377
Peptides
Separation of peptides by multimode chromatography on a column packed with vinyl alcohol copolymer gel 71
Peptides
Isolation and analysis of peptidic fragments of α-gliadin using reversed-phase HPLC 429
Phenazone
Comparison of solid-phase extraction and liquid–liquid extraction for HPLC analysis of antipyrine in plasma 324
Phosphatases
HPLC determination of acid and alkaline phosphatases in urine 196
Phospholipids
Highly sensitive determination and characterization of intact cellular ester-linked phospholipids using LC–plasma spray MS 31
Phospholipids
Behaviour of plasmalogens during HPLC on a silica column with a mobile phase containing phosphoric acid 411
Plasmalogens
Behaviour of plasmalogens during HPLC on a silica column with a mobile phase containing phosphoric acid 411
Primaquine
Review. Detection and determination of antimalarial drugs and their metabolites in body fluids 1
Proguanil
Review. Detection and determination of an-

timalarial drugs and their metabolites in body fluids 1
Propofol
Direct plasma injection using internal surface reversed-phase HPLC: feasibility study using propofol as a model compound 271
Pseudouridine
Determination of age dependency of the creatinine-related pseudouridine excretion in children's urine by HPLC 202
Purine nucleoside phosphorylase
Determination of adenosine deaminase and purine nucleoside phosphorylase activities using HPLC and radiochromatographic methods 439
Purines
Optimization of the ion-pair HPLC separation of purine derivatives in erythrocytes, thymocytes and liver mitochondria 447
Pyridoxal-5-phosphate
Sensitive assay for determination of pyridoxal-5-phosphate in enzymes using HPLC after derivatization with cyanide 209
Pyrimethamine
Review. Detection and determination of antimalarial drugs and their metabolites in body fluids 1
Pyrimidine bases
Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 111
Pyrimidine nucleosides
Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 111
Quinine
Review. Detection and determination of antimalarial drugs and their metabolites in body fluids 1
Retinol
GC of retinol and α-tocopherol without derivatization 21
Sphingolipids
Behaviour of plasmalogens during HPLC on a silica column with a mobile phase containing phosphoric acid 411
Sulphadoxine
Review. Detection and determination of antimalarial drugs and their metabolites in body fluids 1
Temazepam
Computer-aided optimisation of drug enantiomer separation in chiral HPLC 377
Δ^9-Tetrahydrocannabinol
Confirmation of cannabis use. III. Simultaneous quantitation of six metabolites of Δ^9-tetrahydrocannabinol in plasma by HPLC with electrochemical detection 247
Thermooptical detectors
Review. Use of laser detectors in capillary LC 337
Thymine
Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 111
α-Tocopherol
GC of retinol and α-tocopherol without derivatization 21
Triglyceride
Profiling of plasma cholesterol ester and triglyceride fatty acids as their methyl esters by capillary GC, preceded by a rapid aminopropyl-silica column chromatographic separation of lipid classes 395
Trimipramine
Simultaneous determination of trimipramine and its demethylated metabolites in plasma by GC–MS 232
Uracil
Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 111
Uridine
Isocratic HPLC method for studying the metabolism of blood plasma pyrimidine nucleosides and bases: concentration and radioactivity measurements 111
Vanillylmandelic acid
Automated analysis of urinary 3-methoxy-4-hydroxymandelic acid using ion-pair chromatography and fluorimetric detection 43
Vinblastine
Determination of vinca alkaloids in plasma and urine using ion-exchange chromatography on silica gel and fluorescence detection 145
Vinburnine
Simultaneous determination of vinburnine and 6-hydroxyvinburnine in human plasma by HPLC 157
Vinca alkaloids
Determination of vinca alkaloids in plasma and urine using ion-exchange chromato-

graphy on silica gel and fluorescence detection 145
Vindesine
Determination of vinca alkaloids in plasma and urine using ion-exchange chromatography on silica gel and fluorescence detection 145
Vinyl alcohol copolymer gel
Separation of peptides by multimode chromatography on a column packed with vinyl alcohol copolymer gel 71
Vitamin A
GC of retinol and α-tocopherol without derivatization 21
Vitamin E
GC of retinol and α-tocopherol without derivatization 21
Vitride reduction
Application of methylamine O-deacylation and Vitride reduction to the analysis of neutral lipids: a TLC evaluation 215